

SCATTERING BY A NIHILITY SPHERE

Akhlesh Lakhtakia¹

CATMAS — Computational & Theoretical Materials Sciences Group

Department of Engineering Science & Mechanics

212 Earth & Engineering Sciences Building

Pennsylvania State University, University Park, PA 16802-6812

ABSTRACT: On interrogation by a plane wave, the back-scattering efficiency of a nihility sphere is identically zero, and its extinction and forward-scattering efficiencies are higher than those of a perfectly conducting sphere.

Keywords: *back-scattering efficiency; extinction efficiency; forward-scattering efficiency; nihility*

1 INTRODUCTION

The concept of nihility as an electromagnetic *medium* has emerged [1] from the rather extraordinary developments on negatively refracting materials during this decade [2, 3]. Whereas the relative permittivity and relative permeability of vacuum are $\epsilon_r = \mu_r = 1$, and those of anti-vacuum are $\epsilon_r = \mu_r = -1$, those of nihility are $\epsilon_r = \mu_r = 0$. The so-called perfect lens of Pendry [4] is made of anti-vacuum [5], and any perfect lens in the present context is required to simulate nihility [5, 6].

Although quite some attention has been devoted to the electromagnetic response characteristics of anti-vacuum (or some approximation thereof), nihility has been neglected in comparison [7, 8]. Reflection and refraction of plane waves due to nihility half-spaces has recently been reported [9], and so has the plane-wave response of nihility slabs [5, 6]. Along the same lines,

¹E-mail: akhlesh@psu.edu. Also affiliated with: Department of Physics, Imperial College, London SW7 2AZ, United Kingdom

this communication focuses on the plane-wave response of a nihility sphere. An $\exp(-i\omega t)$ time-dependence is implicit in the following sections.

2 THEORY

Consider the spherical region $r < a$ occupied by nihility, whereas the region $r > a$ is vacuous. Without loss of generality, the incident plane wave is taken to be linearly polarized and traveling along the $+z$ axis; thus,

$$\mathbf{E}_{inc}(\mathbf{r}) = E_0 \hat{\mathbf{u}}_x \exp(ik_0 z), \quad (1)$$

where E_0 is the amplitude, k_0 is the free-space wavenumber, and $\hat{\mathbf{u}}_x$ is the unit vector parallel to the $+x$ axis.

As is commonplace, the incident plane wave is represented in terms of vector spherical harmonics $\mathbf{M}_{\sigma mn}^{(j)}(\mathbf{w})$ and $\mathbf{N}_{\sigma mn}^{(j)}(\mathbf{w})$ [10, 11] as follows:

$$\mathbf{E}_{inc}(\mathbf{r}) = E_0 \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} \left[\mathbf{M}_{o1n}^{(1)}(k_0 \mathbf{r}) - i \mathbf{N}_{e1n}^{(1)}(k_0 \mathbf{r}) \right]. \quad (2)$$

The scattered field is also stated in terms of vector spherical harmonics as [11]

$$\mathbf{E}_{sc}(\mathbf{r}) = E_0 \sum_{n=1}^{\infty} i^n \frac{2n+1}{n(n+1)} \left[i a_n \mathbf{N}_{e1n}^{(3)}(k_0 \mathbf{r}) - b_n \mathbf{M}_{o1n}^{(3)}(k_0 \mathbf{r}) \right], \quad r \geq a, \quad (3)$$

where

$$a_n = \frac{\epsilon_r j_n(N\xi) \psi_n^{(1)}(\xi) - j_n(\xi) \psi_n^{(1)}(N\xi)}{\epsilon_r j_n(N\xi) \psi_n^{(3)}(\xi) - h_n^{(1)}(\xi) \psi_n^{(1)}(N\xi)} \quad (4)$$

and

$$b_n = \frac{\mu_r j_n(N\xi) \psi_n^{(1)}(\xi) - j_n(\xi) \psi_n^{(1)}(N\xi)}{\mu_r j_n(N\xi) \psi_n^{(3)}(\xi) - h_n^{(1)}(\xi) \psi_n^{(1)}(N\xi)}; \quad (5)$$

$j_n(\xi)$ and $h_n^{(1)}(\xi)$ are the spherical Bessel function and the spherical Hankel function of the first kind;

$$\psi_n^{(1)}(w) = \frac{d}{dw} [w j_n(w)] \quad (6)$$

and

$$\psi_n^{(3)}(w) = \frac{d}{dw} [w h_n^{(1)}(w)]; \quad (7)$$

$N = \sqrt{\epsilon_r \mu_r}$ and $\xi = k_0 a$; and ϵ_r and μ_r are, respectively, the relative permittivity and the relative permeability of the matter occupying the region $r < a$.

Taking the limits $\epsilon_r \rightarrow 0$ and $\mu_r \rightarrow 0$ for the scattering medium (i.e., nihility), we obtain

$$a_n = b_n = \frac{j_n(\xi)}{h_n^{(1)}(\xi)}. \quad (8)$$

The equality of the coefficients $a_n = b_n \forall n$ is remarkable, and possibly unique to nihility spheres.

3 DISCUSSION

Figure 1 contains a plot of the extinction efficiency

$$Q_{ext} = \frac{2}{\xi^2} \sum_{n=1}^{\infty} (2n+1) \Re(a_n + b_n) \quad (9)$$

of the nihility sphere as a function of its normalized radius ξ . The overall profile is similar to that for a perfectly conducting sphere [13], which is also shown in the same figure, but extinction by the nihility sphere is larger than that by a perfectly conducting sphere. Furthermore, the peak extinction by a nihility sphere occurs at a larger value of ξ (≈ 2.981) than by a perfectly conducting sphere ($\xi \approx 1.209$).

Calculations associated with (4) and (5) show that no electromagnetic field exists inside nihility spheres. Hence, there is no absorption, and the extinction efficiency equals the total scattering efficiency [11]

$$Q_{sca} = \frac{2}{\xi^2} \sum_{n=1}^{\infty} (2n+1) (|a_n|^2 + |b_n|^2). \quad (10)$$

The forward-scattering efficiency

$$Q_{forw} = \frac{1}{\xi^2} \left| \sum_{n=1}^{\infty} (2n+1) (a_n + b_n) \right|^2 \quad (11)$$

of a nihility sphere is plotted in Figure 2 as a function of ξ , and compared with that of a perfectly conducting sphere. That of the nihility sphere is higher.

The most interesting feature of the plane-wave response of a nihility sphere is its back-scattering efficiency [12]

$$Q_{back} = \frac{1}{\xi^2} \left| \sum_{n=1}^{\infty} (-)^n (2n+1) (b_n - a_n) \right|^2. \quad (12)$$

By virtue of (8),

$$Q_{back} \equiv 0 \quad (13)$$

for a nihility sphere; of course, $Q_{back} \neq 0$ for perfectly conducting spheres [12, 13].

Equation (13) is a remarkable result, because it implies that the probability of detection of a nihility sphere by a monostatic radar system is very low. This result would not change even if the ambient (isotropic) medium were to have relative permittivity and relative permeability other than unity. This conclusion can be understood by realizing that nihility is impedance-matched to any isotropic, achiral, dielectric-magnetic medium [14].

References

- [1] A. Lakhtakia, An electromagnetic trinity from “negative permittivity” and “negative permeability”, *Int J Infrared Millim Waves* 22 (2001) 1731–1734; *ibid* 23 (2002) 813–818.
- [2] A. Lakhtakia, M.W. McCall and W.S. Weiglhofer, Brief overview of recent developments on negative phase-velocity mediums (alias left-handed materials), *AEÜ Int J Electron Commun* 56 (2002) 407–410.
- [3] S.A. Ramakrishna, Physics of negative refractive index materials, *Rep Prog Phys* 68 (2005) 449–521.
- [4] J.B. Pendry, Negative refraction makes a perfect lens, *Phys Rev Lett* 85 (2001) 3966–3969.
- [5] A. Lakhtakia, On perfect lenses and nihility, *Int J Infrared Millim Waves* 23 (2002) 339–343.
- [6] A. Lakhtakia and J.A. Sherwin, Orthorhombic materials and perfect lenses, *Int J Infrared Millim Waves* 24 (2003) 19–23.

- [7] S. Tretyakov *et al.*, Waves and energy in chiral nihility, J Electromagn Waves Appl 17 (2003) 695–706.
- [8] R.W. Ziolkowski, Propagation in and scattering from a matched metamaterial having a zero index of refraction, Phys Rev E 70 (2004) 046608.
- [9] A. Lakhtakia and T.G. Mackay, Fresnel coefficients for a permittivity–permeability phase space encompassing vacuum, anti–vacuum, and nihility, Microwave Opt Technol Lett 48 (2006) xxx–xxx.(Issue No. 2)
- [10] J.A. Stratton, Electromagnetic theory, McGraw–Hill, New York, NY, USA, 1941, pp. 414–420.
- [11] C.F. Bohren and D.R. Huffman, Absorption and scattering of light by small particles, Wiley, New York, NY, USA, 1983, ch. 4.
- [12] J. van Bladel, Electromagnetic fields, Hemisphere, Washington, DC, USA, 1985, pp. 264–265.
- [13] V.V. Varadan, A. Lakhtakia and V.K. Varadan, Field representations and introduction to scattering, North–Holland, Amsterdam, The Netherlands, 1991, p. 245.
- [14] P.L.E. Uslenghi, High frequency backscattering from a coated sphere, Alta Freq 34 (1965) 785–788.




Figure 1: Extinction efficiency Q_{ext} of a sphere as a function of its normalized radius $\xi = k_0 a$.
Top: nihility sphere. Bottom: perfectly conducting sphere.




Figure 2: Forward-scattering efficiency Q_{forw} of a sphere as a function of its normalized radius $\xi = k_0 a$. Top: nihility sphere. Bottom: perfectly conducting sphere.